
***Caloneis parva* Levkov & D.M.Williams, nom. nov., a new name for *Caloneis minuta* Levkov & D.M.Williams**

Zlatko Levkov, *Institute of Biology, Faculty of Natural Sciences, Gazi Baba bb, 1000 Skopje, Republic of Macedonia* (corresponding author: zlevkov@iunona.pmf.ukim.edu.mk)

David M. Williams, *Department of Life Sciences, Natural History Museum, Cromwell Road, London, SW7 5BD, UK*

“*Navicula ventricosa* var. *minuta*” first appeared in Cleve & Möller (1879, no. 188) as an invalid *nomen*, but was later validly published as *Navicula ventricosa* f. *minuta* Grunow (in Van Heurck 1880: pl. 12, fig. 26). It was later provided with a description in Van Heurck (1885: 103). Since then it has been recognised as either a variety or form, but most authors agree that it belongs in the genus *Caloneis* Cleve (1894: 46). Cleve, who first described *Caloneis*, also made the first transfer, placing *Navicula ventricosa* f. *minuta* as a variety of *Caloneis silicula*: *Caloneis silicula* var. *minuta* (Grunow) Cleve (Cleve 1894: 52). It was later transferred to *Caloneis ventricosa* var. *minuta* (Grunow) F.W. Mills (Mills 1934: 1177), as the epithet *ventricosa* is the earlier name – Ruth Patrick later made the same, but superfluous, transfer *Caloneis ventricosa* var. *minuta* (Grunow) R.M.Patrick (Patrick & Reimer 1966: 584). Eventually, Ohtsuka and Fujita (2001) made a case for recognising *Caloneis silicula* [*ventricosa*] var. *minuta* as a species and proposed the appropriate change of status to *Caloneis minuta* (Grunow) Ohtsuka & Fujita (Ohtsuka & Fujita 2001: 101).

Regardless of its current taxonomic status, *Caloneis minuta* (Grunow) Ohtsuka & Fujita renders *Caloneis minuta* Levkov & D.M.Williams (2014: 153) an illegitimate later homonym and hence the latter requires a new name:

***Caloneis parva* Levkov & D.M.Williams nomen novum pro *Caloneis minuta* Levkov & D.M.Williams *Nova Hedwigia*, Beihefte 143: 153, 2014, nom. illeg.; non *Caloneis minuta* (Grunow) Ohtsuka & Fujita *Nova Hedwigia* 73(1/2): 101, 2001.**

Cleve, P.T. (1894). Synopsis of the Naviculoid Diatoms. Part I. *Kongliga Svenska-Vetenskaps Akademiens Handlingar* 26(2): 1-194.

Cleve, P.T. & Möller, J.D. (1879). *Diatoms*. Part IV, nos 169-216. Esatas Edquisti Boktryckeri, Upsala.

Levkov, Z. & Williams, DM. (2014). Observations on *Caloneis* Cleve (Bacillariophyceae) species from the ancient lakes Ohrid and Prespa. *Nova Hedwigia*, Beihefte 143: 141-158.

Mills, F.W. (1934). *An index to the genera and species of the Diatomaceae and their synonyms*, 1816-1932. Part 16 (Na-Ni), pp. 1162-1240. Wheldon & Wesley, Limited. London.

Ohtsuka, T. & Fujita, Y. (2001). The diatom flora and its seasonal changes in a paddy field in Central Japan. *Nova Hedwigia* 73(1/2): 97-128.

Patrick, R.M. & Reimer, C.W. (1966). The Diatoms of the United States exclusive of Alaska and Hawaii. Volume 1. Fragilariaeae, Eunotoniaceae, Achnanthaceae, Naviculaceae. *Monographs of the Academy of Natural Sciences of Philadelphia* 13: 1-688.

Van Heurck, H. (1880). *Synopsis des Diatomées de Belgique*. Atlas, pls 1-30. Ducaju & Cie, Anvers.

Van Heurck, H. (1885). *Synopsis des Diatomées de Belgique*. Texte. Martin Brouwers & Co., Anvers.