
The reinstatement of *Hooperia* J.Agardh (*Lomentariaceae*, Rhodophyta) to accommodate *Lomentaria divaricata* (Durant) M.J.Wynne

Michael J. Wynne, *University of Michigan Herbarium, Ann Arbor, Michigan 48108, USA*
(correspondence: mwynne@umich.edu)

Craig W. Schneider, *Department of Biology, Trinity College, Hartford, Connecticut 06106, USA*

Gary W. Saunders, *Centre for Environmental & Molecular Algal Research, Department of Biology, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada*

Multigene analyses of the Rhodymeniales (Rhodophyta) revealed that only two of the six included *Lomentaria* species joined the generitype *L. articulata* (Hudson) Lyngbye in a monophyletic clade viz. *L. clavellosa* (Turner) Gaillon and *L. orcadensis* (Harvey) Collins ex W.R.Taylor (Filloromo & Saunders, 2016). Having resolved in a separate clade as sister to the genus *Ceratodictyon* Zanardini, 1878, *Lomentaria catenata* Harvey was formally transferred to the new genus *Fusuitsunagia* Filloromo & G.W.Saunders, 2016. By contrast, *Lomentaria hakodatensis* Yendo was fully resolved as sister to *Binghamiopsis* I.K.Lee, J.A.West & Hommersand, 1988, but was not subjected to formal taxonomic proposals pending further study (Filloromo & Saunders, 2016). Similarly, *L. divaricata* (Durant) M.J.Wynne resolved as sister to *Stirnia* M.J.Wynne, 2001, but Filloromo & Saunders (2016) chose to refrain from placing *L. divaricata* in the available genus *Hooperia* J.Agardh, 1896 pending future genetic work. Here we propose reinstatement of *Hooperia* J.Agardh to accommodate *L. divaricata*, thus removing its present uncertain nomenclatural placement. *Hooperia* had previously been considered a heterotypic synonym of the earlier genus *Lomentaria* Lyngbye, 1819 (Kylin, 1956).

The type species of *Hooperia* is *H. baileyana* (Harvey) J.Agardh (Agardh 1896), which has *Chylocladia baileyana* Harvey (1853) as its basionym. Wynne (2013) demonstrated that *Chrysomenia divaricata* Durant (Durant 1850) is an older taxonomic synonym of *L. baileyana* (Harvey) Farlow, and so he proposed the binomial *Lomentaria divaricata* (Durant) M.J.Wynne. The following nomenclatural change is proposed:

***Hooperia divaricata* (Durant) M.J.Wynne, C.W.Schneider & G.W.Saunders, comb. nov.**

Basionym: *Chrysomenia divaricata* Durant, *Alg. New York*: 23. 1850.

Homotypic synonym:

Lomentaria divaricata (Durant) M.J.Wynne 2013: 114.

Heterotypic synonyms:

Lomentaria baileyana (Harvey) Farlow 1876: 698.

Chylocladia baileyana Harvey 1853: 185.

Chylocladia divaricata Harvey 1853: 185-186.

Chondrothamnion divaricatum Bailey ex Kützing 1865: 30, pl. 83.

J. Agardh's (1896) name *Hooperia* eponymized British-born American John Hooper (1802–1869), who was greatly influenced by the 1849–1850 visit of William H. Harvey to the United States to self-publish one of the first works on the macroalgae of North America (Hooper 1850).

Agardh, J.G. (1896). *Analecta algologica, Continuatio III. Lunds Universitets Års-Skrift, Andra Afdelningen, Kongl. Fysiografiska Sällskapets i Lund Handlingar* 32(2): 1-140, 1-8 [index], 1 plate.

Durant, C.F. (1850). *Algae and corallines of the Bay & Harbor of New York, illustrated with natural types.* pp. 1-43 and specimens. New York: George P. Putnam Printed by Narine & Co.

Farlow, W.G. (1876). List of the marine algae of the United States. *Report of the United States Fish Commissioner* 1873/1875: 691-717, [1].

-
- Filloramo, G.V. & Saunders, G.W. (2016). Application of multigene phylogenetics and site-stripping to resolve intraordinal relationships in the Rhodymeniales (Rhodophyta). *Journal of Phycology* 52: 339-355.
- Harvey, W.H. (1853). *Nereis boreali-americana*; or, contributions towards a history of the marine algae of the Atlantic and Pacific coasts of North America. Part II. Rhodospermeae. *Smithsonian Contributions to Knowledge* 5: [i-ii], [1]-258, pls XIII-XXXVI.
- Hooper, J. (1850). *Introduction to algology*; with a catalogue of American algae, or sea-weeds, according to the latest classification of Prof. Harvey. pp. [1]-34. Brooklyn: Published by John Hooper.
- Kützing, F.T. (1865). *Tabulae phycologicae*; oder, Abbildungen der Tange. Vol. XV pp. [i-iii], 1-36, 100 pls. Nordhausen: Gedruckt auf kosten des Verfassers (in commission bei W. Köhne).
- Kylin, H. (1956). *Die Gattungen der Rhodophyceen*. pp. i-xv, 1-673, 458 figs. Lund: C.W.K. Gleerups.
- Lee, I.K., West, J.A. & Hommersand, M.H. (1988). *Binghamiopsis caespitosa* gen. et sp. nov. (Lomentariaceae, Rhodophyceae) from the eastern Pacific. *Korean Journal of Phycology* 3: 1-13, 48 figs.
- Lyngbye, H.C. (1819). *Tentamen hydrophytologiae danicae continens omnia hydrophyta cryptogama Daniae, Holsatiae, Faeroae, Islandiae, Groenlandiae hucusque cognita, systematicae disposita, descripta et iconibus illustrata, adjectis simul speciebus norvegicis*. pp. [i]-xxxii, [1]-248, 70 pls. Hafniae [Copenhagen]: typis Schultzianis, in commissis Librariae Gyldendaliae.
- Wynne, M.J. (2001). *Stirnia prolifera* gen. et sp. nov. (Rhodymeniales, Rhodophyta) from the Sultanate of Oman. *Botanica Marina* 44: 163-169, 12 figs.
- Wynne, M.J. (2013). An older name for *Lomentaria baileyana* (Harvey) Farlow (Rhodomiales, Rhodophyta). *Brittonia* 54: 113-117, 1 fig.
- Zanardini, G. (1878). Phyceae papuanae novae vel minus cognitae a cl. O. Beccari in itinere ad Novam Guineam annis 1872-75 collectae. *Nuovo Giornale Botanico Italiano* 10: 34-40.